
## Do you know your Christmas tree species?



Spruce needles are square & sharp



The needles of **pine trees** are fascicled into packets or bundles. The number of needles in each bundle can vary from species, but Lodgepole pines in our area have 2 needles per bundle. These needles are 3 to 9 cm long and are often twisted.



Lodgepole pine trees have stiff ovoid cones from 3 to 6 cm long that grow scattered on the branches. The cones remain on the tree for several years and often look lopsided.





Fir trees have needles with rounded tips which make them friendly to touch. They are also flat in cross section. The needles arise solitary from the stem (not in bundles) and grow from 2 to 3 cm long. The most common fir tree in our area is the Subalpine fir.



Subalpine fir cones are 6 to 10 cm long, grow upward, and are produced toward the top of the tree.

Because fir cones fall apart after maturation, you will rarely find a tree with cones at Christmas time, if you do, they will likely fall apart before you get home.





The needles of **spruce trees** are square in cross section. You can feel this as you roll a needle between your fingertips. Spruce needles are not friendly, they have sharp tips. Needles of the most common spruce native to our area, Engelmann spruce, grow from 1 to 3 cm long.



The cones of Engelmann spruce are light brown in color, are 4 to 7 cm long, and remain on the tree for up to a year after maturing. They are produced toward the top of the tree and hang down from the branches.



## **Other Interesting Christmas tree facts**

A coniferous tree is a 'cone bearing tree'.

Coniferous trees do not produce flowers or fruit
but produce 'naked' seeds inside of cones.

Conifers are referred to as 'evergreen' because their needle-like or scale-like leaves are not shed individually from the tree each year.

Lodgepole pine dwarf mistletoe is a locally native species of dwarf mistletoe. Mistletoes are obligate parasitic plants, anchoring themselves into their host to extract water and nutrients. In the Uinta Mountains, this species can grow on the stems and branches of Lodgepole pine. The leaves of this species of mistletoe are much reduced and scale-like, matching the yellowing to olive green color of the stems. The small, clustered stems end in petal-less flowers. Most of the parasite, however, is beneath the bark causing weaknesses in the host tree.



## The 2010 US Capitol Christmas tree was a 67 foot Engelmann spruce from the Bridger-Teton National Forest in Wyoming.



Douglas fir is another coniferous species that grows in SW Wyoming and the Uinta Mountains. This species is not considered a 'true fir'. Douglas fir needles are flat in cross section but differ from those of the Subalpine fir because the tips that tend to be more blunt than rounded. The needles also narrow slightly before the joining to the twig. Douglas firs are most easily recognized by their cones which have snake-tongue like bracts that stick out between the scales.

